CONTENTS

Potential energy hypersurface determination for non-rigid molecules. II. Single, double internal rotation, and wagging vibration modes
Y.G. Smeyers, M.N. Bellido and A. Niño (Madrid, Spain) 1
A topological characterization for simple molecular surfaces
G.A. Arteca and P. Mezey (Saskatoon, Canada) 11
Resonances in atomic and molecular systems: some simple approaches
R. Lefebvre (Orsay, France) 17
From atom to cosmos
A. Julg (Marseille, France) 33
The Kohn-Sham formalism: a critique and an extension
E.V. Ludeña (Caracas, Venezuela) 39
The formulation and use of density functional theory
J. Keller (México, D.F. México) 51
Double many-body expansion of molecular potential energy functions and the role of long-range forces in the rates of chemical reactions
A.J.C. Varandas (Coimbra, Portugal) 59
An alternative method to assess the quality of Hartree-Fock-Roothaan wave functions
M.I. Menendez, R. Lopez, T.L. Sordo and J.A. Sordo (Oviedo, España)... 75
Natural orbitals in the CIPSI algorithm
G. Berthier (Paris, France), A. Daoudi and J.P. Flament (Palaiseau, France) 81
Use of a matricial form of the T.D.H.F. equations for the calculation of the hyperpolarisability tensor of a particular model
F. Smet and A. Van Groenendael (Villeneuve d'Ascq, France) 87
Gauge-Dependent ket calculations of dynamical polarizabilities of small molecules. Method and application to H_2O
G. Berthier, J.P. Flament and M. Rerat (Palaiseau, France) 97
Spin-Orbit coupling calculations using finite perturbation theory
R. Faiez and M.F. Ruiz Lopez (Vandoeuvre-les-Nancy, France) 103
Full finite perturbation calculations of NMR screening constants and higher order effects
A. Boucekkine, G. Boucekkine-Yaker, M. Nait Achour (Alger, Algeria)
G. Berthier (Paris, France) 109
A pairing theorem for localized molecular orbitals
J.C. Paniagua and A. Moyano (Barcelona, Spain) 113
A variable-β procedure for the calculation of resonance integrals in annulenes through a non-linear formulation of the Hueckel model
J. Marañon and H. Grinberg (La Plata, Argentina) 119
Chimiste: A quantum chemistry computer program
M.T.C. Martins Costa, D. Rinaldi and J.L. Rivail (Vandoeuvre-lès Nancy, France) 125
Computer modelling of protein structures
S. Fraga, R.S. Hodges, J.M.R. Parker, E. San-Fabian and A. Vivier Bunge (Edmonton, Canada) 129
The use of "ab initio" net charges to improve Fraga's atom-atom pair potential for molecular association
F. Torrens, A.M. Sánchez-de-Merás and J. Sánchez-Marín (Burjassot, Spain) 135
Determination of charge densities in molecules containing third period elements. Some critical remarks
Y.G. Smeyers, J.J. Rández, F.J. Rández, M.D. Haro-Ruiz and A. Hernández-Laguna (Madrid, Spain) 141
Ionization potentials of CH_3I: Ab-initio calculations including spin-orbit coupling
M. Tadjeddine, J.P. Flament (Palaiseau, France), C. Teichteil (Toulouse, France) 147

Theoretical computation of the electronic affinity of the BO and BO_2 molecules
F. Mota, J.J. Novoa and A.C. Ramirez (Barcelona, Spain) 153
Approximate relativistic effect on Hartree-Fock type methods
G. Delgado-Barrio, J. Fernández Rico and A. Varadé (Madrid, Spain).. 159
Core polarization corrections on the relativistic core effective potentials formalism
L. Fernández Pacios, M. Fernández Nuñez and P.C. Gomez Calzada (Madrid, Spain) 163
Analysis of the spatial electronic correlation introduced by the omega approach in an atomic system. Application to beryllium
Y.G. Smeyers, G. Delgado-Barrio (Madrid, Spain), L. Doreste-Suarez, J.M. Martín-Gonzalez and E. Ruiz-Egea (Las Palmas de Gran Canaria, Spain) 169
Analysis of the omega function for a molecular system. Application to lithium hydride
L. Doreste-Suarez, J.M. Martín-González (Las Palmas de Gran Canaria, Spain), Y.G. Smeyers and G. Delgado-Barrio (Madrid, Spain) 173
Electronic potentials of alkali hydrides
A. Pardo, J.J. Camacho, J.M.L. Poyato, E. Martín and D. Reyman (Madrid, Spain) 181
A realistic HFACE potential function for $Kr_2(X\,^1\Sigma_g^+)$ from spectroscopic and thermophysical data
J. Brandao, J. Dias da Silva and A.J.C. Varandas (Coimbra, Portugal). 187
Comparison of correlation effects from CI calculations including polarization or eccentric basis functions: BH, CH^+, BeH^- as test cases
G. Boucekkine-Yaker, A. Boucekkine (Alger, Algeria), J.P. Flament (Palaiseau, France),G. Berthier (Paris, France) 193
A reduced basis approach to valence only molecular calculations. Applications to CaO and ScO molecules
S. Polezzo and P. Fantucci (Milano, Italy) 199
A study of simple ionic crystals by a cluster-model approach. Application to the NaF
J.M. Recio and L. Pueyo (Oviedo, Spain) 209
The ab initio perturbed-ion model: A novel approach to the study of crystalline effects on atoms and ions. Application to the NaF
V. Luaña and L. Pueyo (Oviedo, Spain) 215
Comparison of the electronic and geometric structure of Li and Na Clusters. An ab initio quantum chemical study
V. Bonacic-Koutecky and J. Koutecky (Berlin, FRG), P. Fantucci (Milano, Italy) 221
Theoretical calculation of the lower electronic states of the $(MnF_6)^{5-}$ ion
J.F. Van der Maelen, L. Pueyo and M. Bermejo (Oviedo,Spain) 229
Electronic structure of closed-shell hexafluorides
M. Bermejo, V. Luaña, J.M. Recio and L. Pueyo (Oviedo, Spain) 235
Theoretical calculation of the electronic structure of the $(FeF_6)^{3-}$ ion
V.M. García Fernández, G. Fernández Rodrigo and L. Pueyo (Oviedo, Spain.) 241
Structure and analysis of the binding energy of the $CuCO^+$ complex: An ab initio study
P.Y. Morgantini and J. Weber (Geneva, Switzerland) 247
Uncertainty in the intramolecular vibrational energy distribution
M. Fernández-Nuñez and M. Martín Reviejo (Madrid, Spain) 253
Vibrational-rotational energy levels of diatomic molecules from the hill determinant method
D.A. Estrin, F.M. Fernández and E.A. Castro (La Plata, Argentina) ... 257
Internal rotation in molecules with two asymmetric rotors. The symmetry of the hamiltonian
P. Gomez, M. Fernández, L. Fernández Pacios (Madrid, Spain) 261

Rotational barriers of some CH2X radicalar compounds
D. Peeters, G. Leroy, M. Matagne (Louvain-la-Neuve, Belgium) 267
Determination of force constants for the conformational analysis of heteroatomic compounds
M. Ruiz-Pastrana, J.M. Hernando and E. Velasco (Valladolid, Spain). 279
An ab initio quantum chemical study of the formaldimine-water-formic acid system
I.I. Badilescu, C. Sandorfy (Montreal, Canada), C. Mijoule (Paris, France) 283
An ab initio and semiempirical approach to the reaction of zinc hydride with formaldehyde
F. Orsini, F. Pelizzoni (Milano, Italy), D.D. Shillady, L.M. Vallarino (Richmond, USA) 289
Theoretical study of the stability of species A-BH_3(A=Li...Ne)
M.L. Forcada, F. Moscardó and E. San-Fabián (Alicante, Spain) 293
A theoretical study of the addition of atomic boron to water
M. Albertí, R. Sayós, M. González, J. Bofill and A. Aguilar (Barcelona, Spain) 301
Construction of theoretical potentials in van der waals systems. An alternative to the polarization-function counterpoise method
M.A. Martín, S. Tolosa, M.A. Aguilar and F.J. Olivares del Valle (Badajoz, Spain) 307
Solute-solvent interactions. Adaptation of the cavity size as a function of the basis set
M.A. Aguilar, M.A. Martín, S. Tolosa and F.J. Olivares del Valle (Badajoz, Spain) 313
Electrostatical solvent effect and acidities of aliphatic alcohols in DMSO solution
N. Rguini, D. Rinaldi and J.L. Rivail (Vandoeuvre-lès-Nancy, France) 319
Energy levels and dynamics of vibrational predissociation of Ne-Cl_2: Approximate quantum mechanical calculations
P. Villarreal, G. Delgado-Barrio, J. Campos-Martinez and O. Roncero (Madrid, Spain) 325
Thermal rate coefficients for the $^{18}O + {}^{16}O_2 \rightarrow {}^{18}O{}^{16}O + {}^{16}O$ reaction based on a single-valued DMBE potential energy surface for ground-state ozone
A.A.C.C. Pais and A.J.C. Varandas (Coimbra, Portugal) 335
SCF calculations of excited vibrational energy levels for normal modes
A. Hidalgo, J. Zuñiga and A. Requena (Murcia, Spain) 339
Perturbative multi-step adiabatic treatment for energy levels of multidimensional coupled systems
J.M. Frances, J. Zuñiga, A. Hidalgo and A. Requena (Murcia, Spain).. 345
Variational HEG calculation of vibration rotation transition moments for diatomic molecules
J. Zuñiga, F.G. Mateo, A. Hidalgo and A. Requena (Murcia, Spain) ... 351
Perturbed morse expansion for triatomic molecules
J. Zuñiga, A. Bastida, A. Hidalgo and A. Requena (Murcia, Spain) ... 357
Theoretical study of Na*(3p) charge cloud orientation in Na+He collisions
P. Salas, P. Wahnon (Madrid, Spain), C. Courbin (Orsay, France) 363
Photodetachment cross section of analytically tractable systems: The extended asympotic model revisited
G.P. Arrighini, C. Guidotti and N. Durante (Pisa, Italy) 369
Monte Carlo and theoretical study of aqueous solutions of ethanol
G. Alagona and A. Tani (Pisa, Italy) 375
Monte Carlo simulation studies of the solvation of ions. 2. Glycine Zwitterion
G. Alagona, C. Ghio (Pisa, Italy), P.A. Kollman (San Francisco, USA). 385

Monte Carlo evaluation of TST rate constants of elementary processes
E. Vilaseca (Barcelona, Spain) 393
Overlap integrals and memory in stationary perturbation iterative processes for the study of pure liquids
L.M. Sesé, M. Fernández (Madrid, Spain) 397
A preliminary hyperspherical study of the collinear C + HF → CH + F reaction on an "ab initio" potential energy surface
X. Gimenez, J.M. Lucas and A. Aguilar (Barcelona, Spain) 403
A theoretical study of the reaction of $Fe^+(^6D)$ with H_2
M. Sodupe, A. Prenafeta, J.M. Lluch, A. Oliva and J. Bertrán (Barcelona, Spain) 409
Acidity and basicity of azoles: Solvent effects
J. Catalan, M. Sánchez-Cabezudo, J.L.G. de Paz and J. Elguero (Madrid, Spain) 415
Electronic aspects of the hydride transfer mechanism. III. Ab-initio analytical gradient studies of the cyclopropenyl-cation/LiH with 4-31G and 3-21+G basis sets
O. Tapia and R. Cardenas (Uppsala, Sweden), J. Andrés, J.M. Aulló (Valencia, Spain) 421
Thermal rearrangements of the homotropylium cation: An MO investigation of some relevant stationary points on the potential energy surface
M. Barzaghi and C. Gatti (Milan, Italy) 431
Protonation of acetamidoxime: An ab initio molecular orbital study
G. La Manna (Palermo, Italy), M. Cignitti and C.E. Notaro (Roma, Italy) 439
Thermolysis of 3-chloro-3-methyldiazirine. A semiempirical approach
F. Cases, F. Mora and J.M. Pérez (Alicante, Spain) 445
2-R-2-oxo-1,3,2-dioxaphosphorian. II. Electronic structure of the ground state and acido-basic properties
Y. Maouche and B. Maouche (Alger, Algeria), M.C. Brianso (Paris, France) 449
The electronic structure and spectra of 3,6-dipyridyl-s-tetrazines
M. Bossa, G.O. Morpurgo and P. Nota (Roma, Italy) 457
A theoretical study of the electronic spectra of some N-substitued-1,8-naphthalimide
A. Pardo, J. Campanario, J.M.L. Poyato, J.J. Camacho, D. Reyman, E. Martín (Madrid, Spain) 463
The conformation of the OH group in the 3,7 dimethyl-3,7 diazabicyclo [3.3.1] nonan-9-OL molecule
M. Fernández Nuñez, L. Fernández Pacios, F. Sueiro and E. Galvez (Madrid, Spain) 469
Theoretical study of the conformations of cis carbene-acetylene transition metal complexes
B. Maouche (Alger, Algeria), Y. Jean, F. Volatron, O. Eisenstein (Orsay, France) 475
MO Studies on β-lactams. I. On the structure and reactivity of azetidin-2-one with split-valence basis sets
E. Sedano, J.M. Ugalde, F.P. Cossío and C. Palomo (Donostia, Spain). 481
Quantum chemical study of electronic structure and receptor binding in opiates
J.S. Gomez-Jeria and P. Sotomayor (Santiago, Chile) 493
CNDO energies and stability of hydrogen-bonded selfassociated purine and oxipurine dimers
T. Leo (Madrid, Spain), D. Escolar (Cadiz, Spain) 487
The chiral molecule
A. Laforgue (Reims, France) 499